

URSZULA BEDNARZ and IWONA WŁOCH

**Note about sequences of extremal
($A, 2B$)-edge coloured trees**

ABSTRACT. In this paper we determine successive extremal trees with respect to the number of all $(A, 2B)$ -edge colourings.

1. Introduction. For concepts not defined here see [4]. Let G be an undirected, connected and simple graph with the vertex set $V(G)$ and the edge set $E(G)$. Then the order (number of vertices) and the size (number of edges) of G is denoted by n and m , respectively. Let $G(m)$ be a graph of size m . Then $P(m)$, $C(m)$, $T(m)$ and $S(m)$ denote a path, a cycle, a tree and a star of size m , respectively.

Let $P(m_1, m - m_1 - 1)$ be a 2-palm of size m , $m \geq 5$ and the diameter 3 with two support vertices $x, y \in V(P(m_1, m - m_1 - 1))$. Suppose that the support vertex x is adjacent to m_1 leaves, then the vertex y is adjacent to $m - m_1 - 1$ leaves.

In a tree, a vertex of degree at least 3 is a *branch vertex*, a vertex of degree 1 is a *leaf*. If a tree has exactly three leaves, then it is named a *tripod*. In other words, every tripod has the unique branch vertex and consequently this branch vertex is the initial vertex of three elementary paths. Let $m \geq 3$, $p \geq 1$, $t \geq 1$ be integers. Then $T(m, p, t)$ denotes a tripod of size m and three paths of length p, t and $m - p - t$ with the branch vertex as the initial vertex of these paths. For convenience a path of length i , $i \geq 1$ we denote shortly by *i -path*.

2010 *Mathematics Subject Classification.* 11B37, 11C20, 15B36, 05C69.

Key words and phrases. Edge colouring, trees, Fibonacci numbers, telephone numbers.

Let b_m be the number of all nonisomorphic tripods of size m . Then it is given by the following recurrence relation $b_m = 1 + b_{m-2} + b_{m-3} - b_{m-5}$, for $m \geq 5$ with initial conditions $b_0 = b_1 = b_2 = 0$, $b_3 = b_4 = 1$, see [9], [10].

The n th Fibonacci number F_n is defined recursively as follows: $F_n = F_{n-1} + F_{n-2}$, $n \geq 2$ with $F_0 = F_1 = 1$.

The telephone numbers $t(n)$ are defined by the recurrence relation $t(n) = t(n-1) + (n-1)t(n-2)$, for $n \geq 2$ with $t(0) = t(1) = 1$.

Fibonacci and telephone numbers have many interesting applications and interpretations also in graphs. Fibonacci numbers have a graph interpretation known as the Merrifield–Simmons index (i.e. the number of all independent sets) of the n -vertex path P_n , see [5], [6], [7, p. 85–86].

Telephone numbers have also a graph interpretation known as the Hosoya index (i.e. the number of all matchings) of the n -vertex complete graph K_n . For details see [8].

In [1] and [2] there was introduced the graph interpretation of Fibonacci numbers and telephone numbers with respect to the special edge colourings of a graph. We recall this definition.

Let $\mathcal{C} = \{A, B\}$ be the set of two colours. A graph G is $(A, 2B)$ -edge coloured if for every maximal B -monochromatic subgraph H of G there is a partition of H into edge disjoint paths of length 2. Clearly $(A, 2B)$ -edge colouring always exists, since we have no restriction on the colour A .

Let \mathcal{L} be a family of all distinct $(A, 2B)$ -edge coloured graphs obtained by $(A, 2B)$ -colouring of a graph G . Then $\mathcal{L} = \{G^{(1)}, G^{(2)}, \dots, G^{(r)}\}$, $r \geq 1$, where $G^{(p)}$, $1 \leq p \leq r$ denotes a graph obtained by $(A, 2B)$ -edge colouring of a graph G . For $(A, 2B)$ -edge coloured graph $G^{(p)}$, $1 \leq p \leq r$ by $\theta(G^{(p)})$ we denote the number of all partitions of B -monochromatic subgraphs of $G^{(p)}$ into edge disjoint paths of length 2. For the explanation, if $G^{(p)}$ is A -monochromatic, then we put $\theta(G^{(p)}) = 1$. The number of all $(A, 2B)$ -edge colourings we define as the graph parameter

$$\sigma_{(A,2B)}(G) = \sum_{p=1}^r \theta(G^{(p)}).$$

The parameter $\sigma_{(A,2B)}(G)$ was determined for paths, cycles and bounded for trees, for details see [1], [2], [3]. In this paper we give sequences of $(A, 2B)$ -extremal trees, i.e. consecutive trees with extremal values of the parameter $\sigma_{(A,2B)}(T(m))$.

2. Main results. In [2], the lower bound and the upper bound of the parameter $\sigma_{(A,2B)}(T(m))$, $m \geq 1$ were given. Moreover, in [1] it was proved that the upper bound is realized by telephone numbers. This result is presented in the following theorem.

Theorem 1 ([1], [2]). *Let $T(m)$ be a tree of size m , $m \geq 1$. Then*

$$F_m \leq \sigma_{(A,2B)}(T(m)) \leq t(m).$$

Moreover, $\sigma_{(A,2B)}(T(m)) = F_m$ for $T(m) \cong P(m)$ and $\sigma_{(A,2B)}(T(m)) = t(m)$ for $T(m) \cong S(m)$.

Next in [1] the following estimations for the parameter $\sigma_{(A,2B)}(T(m, p, t))$ in the class of tripods were proved:

Theorem 2 ([1]). *Let $m \geq 3$ be an integer. If $T(m) \not\cong P(m)$ and $T(m) \not\cong T(m, p, t)$ for all $p \geq 1$, $t \geq 1$, then*

$$\sigma_{(A,2B)}(P(m)) \leq \sigma_{(A,2B)}(T(m, p, t)) \leq \sigma_{(A,2B)}(T(m)).$$

From the above theorems it is clear that a path $P(m)$ is the extremal tree achieving the minimum value of $\sigma_{(A,2B)}(T(m))$. Moreover, if we want to find the next successive smallest trees with respect to the parameter $\sigma_{(A,2B)}(T(m))$ we have to study the whole class of tripods. The maximum and minimum value of $\sigma_{(A,2B)}(T(m, p, t))$ were established in [3].

Theorem 3 ([3]). *Let $m \geq 4$, $p \geq 1$, $t \geq 1$ be integers. Then*

$$F_{m-1} + 2F_{m-3} \leq \sigma_{(A,2B)}(T(m, p, t)) \leq 2F_{m-1}.$$

Moreover, $\sigma_{(A,2B)}(T(m, p, t)) = 2F_{m-1}$ if $T(m, p, t) \cong T(m, 1, 1)$ and $\sigma_{(A,2B)}(T(m, p, t)) = F_{m-1} + 2F_{m-3}$ if $T(m, p, t) \cong T(m, 2, 2)$.

From Theorem 2 and Theorem 3 we can deduce that the tripod $T(m, 2, 2)$ is the second smallest tree with respect to the $\sigma_{(A,2B)}(T(m))$. In [1] there was found the second minimal tripod $T(m, 4, 2)$ with respect to the parameter $\sigma_{(A,2B)}(T(m))$ which is also, by Theorem 2, the third smallest in the class of trees. From Theorem 3 it is obvious that the tripod $T(m, 1, 1)$ is the largest in the class of tripods with respect to $\sigma_{(A,2B)}(T(m, p, t))$. If we investigate the whole class of tripods, we obtain the sequence of successive extremal tripods from the minimal $T(m, 2, 2)$ to the maximal $T(m, 1, 1)$.

Let $T(m, p, t)$ be an arbitrary tripod, where $m \geq 4$, $p \geq 1$, $t \geq 1$. For $t = 2, 4, \dots, \lfloor \frac{m}{3} \rfloor, \dots, 3, 1$ we obtain the successive smallest tripods with respect to the parameter $\sigma_{(A,2B)}(T(m))$. The integer t assumes the consecutive even numbers from 2 to $\lfloor \frac{m}{3} \rfloor$, if $\lfloor \frac{m}{3} \rfloor$ is even or to $\lfloor \frac{m}{3} \rfloor - 1$, if $\lfloor \frac{m}{3} \rfloor$ is odd. Then t assumes the consecutive odd numbers from $\lfloor \frac{m}{3} \rfloor$ or $\lfloor \frac{m}{3} \rfloor - 1$ to 1. In other words, the tripod $T(m, p, 2)$ always achieves the smallest value of $\sigma_{(A,2B)}(T(m, p, t))$ and the tripod $T(m, p, 1)$ always achieves the largest value of this parameter.

Moreover, for each value of t we can construct a sequence of extremal tripods with respect to the integer p .

If $t = 1$, then the successive smallest tripods are given respectively for $p = 2, 4, \dots, \lfloor \frac{m-1}{2} \rfloor, \dots, 3, 1$.

If $t = 2$, then $p = 2, 4, \dots, \lfloor \frac{m-2}{2} \rfloor, \dots, 5, 3$.

If $t = 3$, then $p = 4, 6, \dots, \lfloor \frac{m-3}{2} \rfloor, \dots, 5, 3$.

If $t = 4$, then $p = 4, 6, \dots, \lfloor \frac{m-4}{2} \rfloor, \dots, 7, 5$.

\vdots

If $t = \lfloor \frac{m}{3} \rfloor$ and $\lfloor \frac{m}{3} \rfloor$ is even, then $p = \lfloor \frac{m}{3} \rfloor, \dots, \lfloor \frac{m - \lfloor \frac{m}{3} \rfloor}{2} \rfloor, \dots, \lfloor \frac{m}{3} \rfloor + 1$,

and if $\lfloor \frac{m}{3} \rfloor$ is odd, then $p = \lfloor \frac{m}{3} \rfloor + 1, \dots, \lfloor \frac{m - \lfloor \frac{m}{3} \rfloor}{2} \rfloor, \dots, \lfloor \frac{m}{3} \rfloor$.

For example if we consider the tripods $T(12, p, t)$ for all possible p, t , then the sequence of successive smallest tripods with respect to the increasing parameter $\sigma_{(A,2B)}(T(m, p, t))$ is as follows

$$T(12, 2, 2), T(12, 4, 2), T(12, 5, 2), T(12, 3, 2), T(12, 4, 4), T(12, 4, 3),$$

$$T(12, 3, 3), T(12, 2, 1), T(12, 4, 1), T(12, 5, 1), T(12, 3, 1), T(12, 1, 1).$$

The above considerations give an answer to the question given in [1].

Let $t_i(m)$, $i = 1, \dots, b_{m+1}$ be the i th minimum tree of size m with respect to the parameter $\sigma_{(A,2B)}(T(m))$. Then

$$t_1(m) \cong P(m), t_2(m) \cong T(m, 2, 2), t_3(m) \cong T(m, 4, 2),$$

$$t_4(m) \cong T(m, 6, 2), \dots, t_{m-1}(m) \cong T(m, 5, 1),$$

$$t_m(m) \cong T(m, 3, 1), t_{m+1}(m) \cong T(m, 1, 1).$$

Now we determine the successive trees with respect to the decreasing parameter $\sigma_{(A,2B)}(T(m))$. To do this we use among others the following lemma.

Lemma 1 ([2]). *Let $G = H \cup T(l) \cup \{e\}$ be a connected graph, where H is a connected graph, $T(l)$ is a tree of size l , $l \geq 1$ and H and $T(l)$ are vertex disjoint. Assume that $e = uv$, where $u \in V(H)$ and $v \in V(T(l))$. Then*

$$(1) \quad \sigma_{(A,2B)}(H \cup P(l) \cup \{e\}) \leq \sigma_{(A,2B)}(G) \leq \sigma_{(A,2B)}(H \cup S(l) \cup \{e\}),$$

where the vertex v is identified with the center of the star $S(l)$. Moreover, the equality holds if $T(l) \cong P(l)$ or $T(l) \cong S(l)$.

Theorem 4. *Let $m \geq 5$, $m_1 \geq 2$ be integers. Then*

$$\sigma_{(A,2B)}(T(m)) \leq \sigma_{(A,2B)}(P(m_1, m - m_1 - 1)) \leq \sigma_{(A,2B)}(S(m)).$$

Proof. The inequality $\sigma_{(A,2B)}(P(m_1, m - m_1 - 1)) \leq \sigma_{(A,2B)}(S(m))$ follows immediately from Theorem 1. Let $T(m) \not\cong S(m)$ and $T(m) \not\cong P(m_1, m - m_1 - 1)$. Then $\text{diam } T(m) \geq 4$. Let $\bar{P} = x - y$ be the path which realizes the diameter $\text{diam } T(m)$. Then $x, y \in V(T(m))$ are leaves. Let $u \in V(T(m))$ be adjacent to the vertex x and the edge $e \in \bar{P}$ be incident with u and e is not incident with a leaf. Then $T(m) = T(m_1) \cup T(m_2) \cup \{e\}$, where $m_1 + m_2 + 1 = m$. Applying Lemma 1, we have

$$\sigma_{(A,2B)}(T(m)) \leq \sigma_{(A,2B)}(S(m_1) \cup S(m_2) \cup \{e\}) = \sigma_{(A,2B)}(P(m_1, m - m_1 - 1)),$$

which ends the proof. \square

Let $T_i(m)$ be the i th maximum tree of size m with respect to the parameter $\sigma_{(A,2B)}(T(m))$.

Theorem 5. *Let $m \geq 5$ be an integer. Then $T_1(m) = S(m)$, $\sigma_{(A,2B)}(S(m)) = t(m)$ and $T_i(m) = P(m-i, i-1)$ for $2 \leq i \leq m - \lceil \frac{m-1}{2} \rceil$.*

Theorem 6. *Let $m \geq 5$, $m_1 \geq 2$ be integers. If $T(m) \not\cong S(m)$ and $T(m) \not\cong P(m_1, m-m_1-1)$, then $\sigma_{(A,2B)}(T(m)) \leq \sigma_{(A,2B)}(P(m_1, m-m_1-2))$.*

Proof. Since $T(m) \not\cong S(m)$ and $T(m) \not\cong P(m_1, m-m_1-1)$ then $\text{diam } T(m) \geq 4$. Let $e \in E(T(m))$ is not incident with a leaf. Such edge there exists because $\text{diam } T(m) \geq 4$. Let $T(m) = T(m_1) \cup \{e\} \cup T(m_2)$, where $m = m_1 + m_2 + 1$. Then $\text{diam } T(m_1) \geq 2$ or $\text{diam } T(m_2) \geq 2$. Suppose without loss of generality that $\text{diam } T(m_2) \geq 2$. Let us consider the following possibilities:

1. $c(e) = A$. Then

$$\sigma_{A(e)}(T(m)) = \sigma_{(A,2B)}T(m_1)\sigma_{(A,2B)}T(m_2).$$

2. $c(e) = 2B$. Then

$$\begin{aligned} \sigma_{2B(e)}(T(m)) &= \sigma_{2B(e)}(T(m_1) \cup \{e\})\sigma_{(A,2B)}(T(m_2)) \\ &\quad + \sigma_{(A,2B)}(T(m_1))\sigma_{2B(e)}(T(m_2) \cup \{e\}). \end{aligned}$$

Therefore $\sigma_{(A,2B)}(T(m)) = \sigma_{A(e)}(T(m)) + \sigma_{2B(e)}(T(m))$. Hence

$$\begin{aligned} \sigma_{(A,2B)}(T(m)) &= \sigma_{(A,2B)}T(m_1)\sigma_{(A,2B)}T(m_2) \\ &\quad + \sigma_{(A,2B)}(T(m_1) \cup \{e\})\sigma_{(A,2B)}(T(m_2)) \\ &\quad + \sigma_{(A,2B)}(T(m_1))\sigma_{(A,2B)}(T(m_2) \cup \{e\}). \end{aligned}$$

Since $\text{diam } T(m_2) \geq 2$ then applying Lemma 1, we have

$$\begin{aligned} \sigma_{(A,2B)}(T(m)) &\leq \sigma_{(A,2B)}(P(m_2-1, 1))\sigma_{(A,2B)}(S(m_1)) \\ &\quad + \sigma_{2B(e)}(P(m_2-2, 1))\sigma_{(A,2B)}(S(m_1)) \\ &\quad + \sigma_{(A,2B)}(P(m_2-1, 1))\sigma_{2B(e)}(P(m_2-1, 1)) \\ &= \sigma_{(A,2B)}(P(m_2-1) \cup S(m_1) \cup \{e\}) \\ &= \sigma_{(A,2B)}(P(m_1, m-m_1-2)) \end{aligned}$$

which ends the proof. \square

Theorem 7. *Let $m \geq 5$ be an integer. Then*

$$\sigma_{(A,2B)}(P(m_1, m-m_1-2)) = t(m_1+1)t(m-m_1-1) + t(m_1)t(m-m_1-2).$$

Proof. Let $e, e' \in E(P(m_1, m-m_1-2))$ be not incident with a leaf. If $c(e) = c(e') = 2B$ and a 2-path $e - e'$ belongs to a partition of $2B$ -monochromatic subgraph into 2-paths, then we have $t(m_1)t(m-m_1-2)$ possibilities in this case. Otherwise the tree $P(m_1, m-m_1-2)$ can be considered as the union of two stars $S(m_1+1)$ and $S(m-m_1-1)$ and the result follows. \square

From the above theorems we have

Corollary 8. *Let $m \geq 5$, $m_1 \geq 2$ be integers, $T(m) \not\cong S(m)$ and $T(m) \not\cong P(m_1, m - m_1 - 1)$. Then*

$$\sigma_{(A,2B)}(T(m)) \leq t(m_1 + 1)t(m - m_1 - 1) + t(m_1)t(m - m_1 - 2).$$

In the same way as for the palm $P(m_1, m - m_1 - 1)$, see [1], we can show the behavior of the parameter $\sigma_{(A,2B)}(P(m_1, m - m_1 - 2))$ after moving an edge adjacent to a support vertex to another support vertex. So we omit the proof.

Lemma 2. *Let $m \geq 6$, $m_1 \geq 2$ be integers and $m_1 \geq m - m_1 - 2$. Then*

$$\sigma_{(A,2B)}(P(m_1 + 1, m - m_1 - 3)) > \sigma_{(A,2B)}(P(m_1, m - m_1 - 2)).$$

Theorem 9. *Let $T(m)$ be a tree of size m , $m \geq 6$, $T(m) \not\cong S(m)$ and $T(m) \not\cong P(m_1, m - m_1 - 1)$ for all $m_1 \geq 2$. Then*

$$\sigma_{(A,2B)}(T(m)) \leq \sigma_{(A,2B)}(P(m - 3, 1)).$$

Proof. Let $T(m)$ be a tree of size m , $m \geq 6$ such that $T(m) \not\cong S(m)$ and $T(m) \not\cong P(m_1, m - m_1 - 1)$ for all $m_1 \geq 2$. Then by Theorem 6, $\sigma_{(A,2B)}(T(m)) \geq \sigma_{(A,2B)}(P(m_1, m - m_1 - 2))$ for all $m \geq 2$. If $m - m_1 - 2 = 1$, then $m_1 = m - 3$ and $P(m_1, m - m_1 - 2) \not\cong P(m - 3, 1)$, so the result follows. Let $m - m_1 - 2 \geq 2$ and without loss of the generality, suppose that $m_1 \geq m - m_1 - 2$. Applying Lemma 2, we obtain

$$\sigma_{(A,2B)}(P(m_1, m - m_1 - 2)) < \sigma_{(A,2B)}(P(m_1 + 1, m - m_1 - 3)).$$

If $m - m_1 - 3 \geq 2$, then we apply Lemma 2 until we obtain the palm $P(m - 3, 1)$, which ends the proof. \square

Let $T_i^*(m)$ be the i th maximum tree of size m in the class of 2-palms $P(m_1, m - m_1 - 2)$ for $m_1 \geq 2$ with respect to the parameter $\sigma_{(A,2B)}(T(m))$. From the above considerations we have

Theorem 10. *Let $m \geq 5$ be an integer. Then $T_i^*(m) = P(m - i - 2, i)$ for $i = 1, 2, \dots, \lceil \frac{m-2}{2} \rceil$.*

REFERENCES

- [1] Bednarz, U., Włoch, I., *Fibonacci and telephone numbers in extremal trees*, Discuss. Math. Graph Theory, doi 10.7151/dmgt.1997, in press.
- [2] Bednarz, U., Włoch, I., Wołowicz-Musiał, M., *Total graph interpretation of numbers of the Fibonacci type*, J. Appl. Math. **2015** (2015), ID 837917, 7 pp.
- [3] Bednarz, U., Bród, D., Szynal-Liana, A., Włoch, I., Wołowicz-Musiał, M., *On Fibonacci numbers in edge coloured trees*, Opuscula Math. **37** (4) (2017), 479–490.
- [4] Diestel, R., *Graph Theory*, Springer-Verlag, Heidelberg, New York, 2005.
- [5] Gutman, I., Wagner, S., *Maxima and minima of the Hosoya index and the Merrifield–Simmons index. A survey of results and techniques*, Acta Appl. Math. **112** (3) (2010), 323–346.

-
- [6] Prodinger, H., Tichy, R. F., *Fibonacci numbers of graphs*, Fibonacci Quart. **20** (1982), 16–21.
- [7] Riordan, J., *Introduction to Combinatorial Analysis*, Dover Publ., Inc., New York, 2002.
- [8] Tichy, R. F., Wagner, S., *Extremal problems for topological indices in combinatorial chemistry*, J. Comput. Biol. **12** (7) (2005), 1004–1013.
- [9] Weisstein, E., *Tripod index entries for linear recurrence with constant coefficients*, MathWorld, Wolfram Web Resource, Mar. 05 2011,
URL <http://mathworld.wolfram.com/Tripod.html>
- [10] *The On-Line Encyclopedia of Integer Sequences*,
URL <https://oeis.org/>

Urszula Bednarz
Faculty of Mathematics
and Applied Physics
Rzeszów University of Technology
al. Powstańców Warszawy 12
35-359 Rzeszów
Poland
e-mail: ubednarz@prz.edu.pl

Iwona Włoch
Faculty of Mathematics
and Applied Physics
Rzeszów University of Technology
al. Powstańców Warszawy 12
35-359 Rzeszów
Poland
e-mail: iwloch@prz.edu.pl

Received March 23, 2017